

The Effect of Harvest Time on Bioactive Compounds of Field-Cultivated *Centella asiatica* (L) Urban

Noverita Sprinse Vinolina¹, Luthfi Siregar¹, Bilter Sirat², Nurhayati³, Enny Adelina⁴, Laila Nazirah⁵

- 1) Agriculture Faculty, Department of Agrotechnology, Universitas Sumatera Utara, North Sumatra, Indonesia
- 2) Agriculture Faculty, Department of Agrotechnology, Universitas Darma Agung, North Sumatra, Indonesia
- 3) Agriculture Faculty, Department of Agrotechnology, Universitas Islam Sumatera Utara, North Sumatra, Indonesia
- 4) Agriculture Faculty, Department of Agrotechnology, Universitas Tadulako, Central Sulawesi, Indonesia
- 5) Agriculture Faculty, Department of Agrotechnology, Universitas Malikussaleh, Aceh, Indonesia

Main author's email, phone number & address:

noveritasitumorang@yahoo.com, (62) 812 6052 933

Agriculture Faculty, Department of Agrotechnology, University of Sumatera Utara Jl. Dr. A. Sofian No.3, Padang Bulan, Medan Baru, Medan City, North Sumatra 20155

Running title: Effect of Harvest Time on Pegagan's Bioactive Compounds

Keywords: *Centella asiatica* (L) Urban, bioactive compounds, field cultivation, harvest time, centellosides

Abstract

Background: Pegagan, *Centella asiatica* (L) Urban is an important traditional herb, and its wild population is currently heavily exploited due to commercial cultivation shortage. **Objectives:** This study evaluated the content of different bioactive compounds in the leaves and roots of Pegagan, *Centella asiatica* (L) Urban cultivated in field condition and harvested at different times. **Materials and Methods:** The contents of centellosides in the leaves and roots of field-cultivated Deli Serdang Pegagan Accession were determined by using ultra fast liquid chromatography system. The levels of bioactive compounds in relation to harvest time were compared with an additional discussion on the dry/wet weight relationship. **Results:** The result showed that asiaticoside and madecasoside increased while asiatic acid decreased over time in both the leaves and roots of Pegagan. However, the leaves have significantly higher amounts of all centellosides compared to the roots in all harvest time treatments. **Conclusion:** This study concludes that a late harvest of Pegagan is best to maximize the production of asiaticoside and madecasoside. Conversely, early harvest is best to the maximum amount of asiatic acid in Pegagan. Farmers are also recommended in mainly harvesting the leaves as it is more economically profitable in terms of lower wet to dry weight ratio and the maximum contents of centelloside compounds.

Keywords: *Centella asiatica* (L) Urban, bioactive compounds, field cultivation, harvest time, centellosides, Pegagan

INTRODUCTION

Centella asiatica or widely known as Gotu Kola in the international market or 'Pegagan' in Indonesia¹ is known for its bioactive compounds and has been used as a medicinal ingredient to treat various illnesses and diseases²⁻⁴. Such benefits offered by this tropical plant has led to its commercial cultivation in some of the dominant producing countries such as China and India^{5,6}. However, the commercial cultivation of Pegagan in Indonesia is lagged behind due to the inconsistent supply of quantity and quality due to environmental conditions, harvest and post-processing⁷. As a result, the increasing national and international demands of Pegagan is primarily supplied from the wild population⁸ resulting in overexploitation of the plant as has happened worldwide⁹.

Previous studies on the content of bioactive compounds in Pegagan such as asiaticoside, madecassoside, and asiatic acid have been focusing on their complex and benefits of biochemical processes¹⁰, their concentration in different parts of the plant¹¹⁻¹³ and general reviews of its growth under controlled conditions^{11,14}. There were limited or non-existent literature or studies, particularly within Indonesian regions, that provide information regarding the effect of harvest time on the concentration of important bioactive compounds of Pegagan cultivated in field condition. Such discrepancy has been highlighted by Ghulamahdi, *et al.*¹⁵ and Vinolina, *et al.*⁷ where the field production of Pegagan varies in terms of quality and quantity throughout the year. Field location, seasonal climate condition as well as accession type contributed to these inconsistencies and needed to be studied further^{10,16}

This recent study determined the content of different bioactive compounds in different parts of field-cultivated Pegagan harvested at different times. The information is important to fill the gap of knowledge regarding the appropriate harvest time to maximize the production quantity and quality of certain bioactive compounds in Pegagan cultivated in field condition.

MATERIALS AND METHODS

Study Location

The study was carried out for nine months from January to September 2014. The experimental field was located at the University of Sisingamangaraja XII, Medan, North Sumatra situated at an altitude of 50 m above sea level.

Seed Production and Field Preparation

This study used Gotu Kola seeds from the accession of Deli Serdang, North Sumatra, Indonesia due to its higher content of centellosides in a previous study by Vinolina and Siregar¹⁷. The mother plants were grown for two and a half months and used as the seeds. The seeds were planted in previously prepared 30 units of 30 cm raised-soil beds or plots sized 1 m x 1 m. Each treatment (harvest time) used 10 plots where four seeds of Pegagan was planted in each plot at 40 cm distance. Randomized Block Design was used to arrange the treatments consisted of different harvest time of 56, 70, and 84 days after planting denoted as U₁, U₂ and U₃, respectively. Each treatment had 10 replicates to determine the effect of harvest time on biomass production of and bioactive compound in Pegagan. Fertilization was conducted at

0 days after planting (DAP) and repeated at 20 and 40 DAP using superphosphate 36 (SP36), Urea, and KCL of 20 g, 30 g, and 22 g, respectively, for each plot. Field maintenance was regularly conducted to maintain the optimum growth of cultivated Pegagan.

Harvest Procedures

The harvest times of cultivated Pegagan as the treatment variable in this current study were set at day 56, 70, and 84 after planting denoted as U_1 , U_2 , and U_3 , respectively. Before harvest, the plots were wetted with water to facilitate easy removal of the plants. Each whole plant was taken out from the soil plots by hand. Cleaned plants were grouped according to the harvest times, and part categories (leaves and roots) then weighed to determine their wet weights. Afterward, all samples were oven-dried for three days at 50 °C and ground to powder for the subsequent bioactive compound analysis.

Bioactive Compounds in Pegagan

The powder sample harvested Pegagan (0.2 g for each sample) was prepared by pouring into a test tube, mixed with 4 ml of 90% methanol solution, shaken for five hours. The supernatant was collected and then evaporated using a waterbath set at 50 °C. The dried supernatant was then mixed with 1 ml of 90% methanol solution (gradient grade for liquid chromatography), filtered using a 0.22 μ M PTFE filter (Whatman membrane filter P/N E252 Buckinghamshire, England) and ready for the subsequent ultra fast liquid chromatography (UFLC) process.

The analyses of bioactive compounds were performed at the Pharmacy Laboratory of the University of North Sumatera. Standard commercial asiaticoside, madecassoside, and asiatic acid were obtained from Sigma Aldrich (St. Louis, MO, USA) and used to determine the retention time of each compound following the modified procedure of Jain and Agrawal¹⁸. The calibration of the UFLC Shimadzu DGU-20A3 followed the procedures described by Mangas, *et al.*¹⁹ and Bonfill, *et al.*¹³. The retention time of asiaticoside, madecasoside and asiatic acid were determined as 18.603, 13.196, and 21.054 min, respectively. The contents of asiaticoside, madecasoside, and asiatic acid were determined by injecting 20 μ l of the filtrates of each sample in the UFLC system with the retention times previously described. The measurement of each centelloside compound was repeated three times, and the values were averaged.

Statistical Analysis

Experimental data were analyzed using F-test in the analysis of variance (ANOVA) and significant differences among the treatments ($P < 0.05$) were determined by Duncan's multiple range test (DMRT) using the SPSS 17.0 software.

RESULTS

Asiaticoside concentration at different harvest times

The concentrations of asiaticoside in leaves (KAA, $\mu\text{g/mL}$) and roots (KAD, $\mu\text{g/mL}$) of Pegagan harvested at a different time (U_1 , U_2 , and U_3) are presented in Figure 1. The figure clearly shows that both Pegagan parts contained significantly different asiaticoside amounts according to the harvest times. The highest concentration of asiaticoside was found in the leaves of Pegagan ($53.55 \mu\text{g/mL}$) harvested at 70 DAP which was more than four times compared to that of the leaves harvested at 56 DAP ($12.52 \mu\text{g/mL}$). The ratio of asiaticoside content in the leaves and roots of U_2 differed greatly compared to that of U_1 and U_3 ($p < 0.05$).

Madecasosside concentration at different harvest times

The concentrations of madecasosside showed significant differences in different parts (leaves and roots) and harvest times (Figure 2) ($p < 0.05$). The leaves consistently had the highest concentration of madecasosside compared with the roots across different harvest times. The highest concentration of madecasosside was found in the leaves of Pegagan harvested at U_3 ($358.18 \mu\text{g/mL}$) of almost four times of the lowest concentration in the leaves harvested at U_2 and more than 17 times of the lowest concentration in the roots harvested at U_2 .

Asiatic Acid concentration at different harvest times

Figure 3 shows that the contents of asiatic acid in the leaves of Pegagan were consistently high compared that of the roots ($p < 0.05$). However, there was a clear decreasing pattern of asiatic acid content in the leaves (KAAD) of Pegagan from the shortest to the longest harvest times. The content of asiatic acid in Pegagan's leaves harvested had decreased from the shortest harvest time (U_1 : $611.6 \mu\text{mL}$) to less than half in Pegagan's leaves harvested at the longest time (U_3 : $294.56 \mu\text{mL}$). Despite the contents of asiatic acid in the roots (KAAA) were relatively stable at different harvest times, the F-test showed significant differences among the treatments with U_3 had the highest asiatic acid, followed by U_1 and the lowest was U_2 ($p < 0.05$).

Dry and wet weight of harvested Pegagan at different times

The wet weight (WW) and dry weight (DW) of Pegagan harvested at U_1 , U_2 , and U_3 and dried in an oven at 50°C for three days is presented in Table 1. The WWs and DWs of leaves and roots of harvested Pegagan at U_3 were the highest and U_1 were the lowest ($p < 0.05$).

DISCUSSION

This current study has found that there is an opposite relationship between the contents of asiaticoside -madecasosside and asiatic acid in both the leaves and roots of Pegagan cultivated at field condition. Asiaticoside - madecasosside content in whole plant of Pegagan increases over time whilst at the same time, the content of asiatic acid decreases. This finding reconfirms the

findings of Kim, *et al.*¹¹ and Bonfill, *et al.*¹³ who had found similar results with Pegagan cultivated under laboratory condition and treated with elicitors to increase the contents of centellosides. However, the level of centellosides in the whole plant of Pegagan found in this study was significantly higher particularly for asiaticoside which was up to five-fold higher compared to that of Bonfill, *et al.*¹³ but relatively similar to that of Kim, *et al.*¹¹. Here we argue that the relatively higher content of centelloside found in this study is due to the twice longer cultivation period used by the two studies. This longer cultivation period had allowed the build-up of the secondary metabolites in the leaves and roots of Pegagan despite no elicitors used. A similar result was also reported by Rafi, *et al.*¹ that harvest time determined how much concentration of centelloside to be accumulated in whole plant Pegagan. However, their study did not distinguish the concentrations of different centellosides in leaves and roots of Pegagan. The field condition in which the Pegagan were cultivated is also argued to support all nutrients required by the plant (Table 2) compared to the more sterile and controlled laboratory condition. This was described by Jamil, *et al.*²⁰ and Rafi, *et al.*¹ as the effect of geographic location and environmental condition which also have the role the production of centellosides in Pegagan.

Furthermore, the patterns of centellosides show a consistent trend that the leaves of Pegagan had a higher content of centellosides compared to the roots across different harvest times at field condition. A laboratory study by Mangas, *et al.*¹⁹ and Kim, *et al.*¹¹ also found that Pegagan mainly synthesizes the secondary metabolites in leaves and an only limited amount in the roots. Similarly, Zainol, *et al.*²¹ had found that leave of Pegagan has the highest concentration of centellosides compared to the root. A detailed study by Gupta, *et al.*¹⁶ also reported a high concentration of different centellosides in a similar order as found in this study. However, they studied the morphometric variation in relation to bioactive compounds of Pegagan leaves sourced from the wild population which is not comparable to this study. There are currently no studies explaining the tendency. However, we suspect that accumulation of the secondary metabolites in the leaves of Pegagan is mainly for defense mechanism to predation and diseases due to its perennial nature. Rosenthal and Berenbaum²² argue that secondary metabolites occur in a large amount in a particular plant part as a defense mechanism to predation and diseases. Such a defense mechanism might explain some woody plants produce a higher amount of secondary metabolites in roots compared in the leaves. For example, cassava produces cyanogenic glucosides in its cassava tubers to deter root borers²³. This might be the case for Pegagan where the higher amount of centellosides in leaves serves as a deterrent for disease, pest, and predators although another study is required to confirm this.

In addition, the roots' dry weight of Pegagan was relatively heavier compared to that of the leaves across different harvest times (Table 1). This implies that it is more economical to harvest the leaves than the roots because leaves are easier to harvest and have less weight compared to the root. It also corresponds to the higher amount of asiaticoside and madecacoside in the leaves that can be synthesized compared to that of the roots^{4,5}.

Future research direction regarding the effect of harvest time on field-cultivated Pegagan can be focused on the addition of elicitors such as Methyl Jasmonate to stimulate the production of centellosides in Pegagan. The research on Methyl Jasmonate to increase

centellosides in Pegagan cultivated under laboratory or controlled condition are many and show a promising result. However, its application in field-cultivated Pegagan has yet to be done.

CONCLUSION

The study revealed that the Pegagan accession of Deli Serdang, Indonesia has asiaticoside and madecasoside in the leaves and roots which increases and asiatic acid which decreases over time as the plant aged. All the three centellosides are mainly found the leaves and very little in the roots across different harvest times suggesting that these secondary metabolites are for defense mechanism from pest, predation, and diseases due to the closeness of aerial parts of Pegagan to the ground.

SIGNIFICANCE STATEMENT

This study discovered different harvest time on Pegagan can be beneficial for maximizing the level of bioactive compounds in commercial field cultivation of this plant. This study presents new findings regarding bioactive compounds level of Pegagan cultivated in field-condition and harvested at different times, which are currently lacking in the literature. This study will help researchers, farmers and biopharmaceutical industries to uncover the critical areas of harvest time settings of field-cultivated Pegagan in order to maximize bioactive compounds in the plant that many researchers and practitioners were not able to explore. Thus, a new theory on how harvest time and environmental factors affect the plant production of bioactive compounds may be arrived at.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the research funding support, grant number 097/K1.1.1/AT.1/2015, from the Ministry of Research, Technology and Higher Education, the Republic of Indonesia.Page Break

REFERENCES

- 1.Rafi, M., Handayani, F., Darusman, L.K., Rohaeti, E., Wahyu, Y., Sulistiyani, Honda, K. and Putri, S.P. 2018. A combination of simultaneous quantification of four triterpenes and fingerprint analysis using HPLC for rapid identification of *Centella asiatica* from its related plants and classification based on cultivation ages. *Ind. Crops Prod.*, 122:93-97. Available from: <https://doi.org/10.1016/j.indcrop.2018.05.062> [Accessed 28 October 2018].
- 2.James, J. and Dubery, I. 2011. Identification and quantification of triterpenoid centelloids in *Centella asiatica* (L.) Urban by densitometric TLC. *JPC-J Planar Chromat.*, 24(1):82-87. Available from: <https://doi.org/10.1556/JPC.24.2011.1.16> [Accessed 12 December 2018].
- 3.Ncube, E.N., Steenkamp, P.A., Madala, N.E. and Dubery, I.A. 2017. Metabolite profiling of the undifferentiated cultured cells and differentiated leaf tissues of *Centella asiatica*. *Plant Cell Tissue Organ Cult.*,129(3):431-443. Available from: <https://doi.org/10.1007/s11240-017-1189-4> [Accessed 3 October 2018].
- 4.Mangas, S., Moyano, E., Hernandez-Vazquez, L. and Bonfill, M. 2009. In: *Plant Secondary Terpenoids, Centella asiatica* (L) Urban: An Updated Approach Terpenoids, Palazón J, Cusidó RM,(Eds.) Fort P.O., Trivandrum-695 023, Kerala, India. Research Signpost, pp: 55-74.
- 5.Ariffin, F., Heong Chew, S., Bhupinder, K., Karim, A.A. and Huda, N. 2011. Antioxidant capacity and phenolic composition of fermented *Centella asiatica* herbal teas. *J. Sci. Food Agric.*,91(15):2731-2739. Available from: <https://doi.org/10.1002/jsfa.4454> [Accessed 11 November 2018].

6. Gallego, A., Ramirez-Estrada, K., Vidal-Limon, H.R., Hidalgo, D., Lalaleo, L., Khan Kayani, W., Cusido, R.M. and Palazon, J. 2014. Biotechnological production of centellosides in cell cultures of *Centella asiatica* (L) Urban. *Eng. Life Sc.*,14(6):633-642. Available from <https://doi.org/10.1002/elsc.201300164> [Accessed 1 January 2019].
7. Vinolina, N.S., Nainggolan, M. and Siregar, R. 2018. Production Enhancement Technology of Pegagan (*Centella asiatica*). *AGRIVITA*,40(2):304-312. Available from: <http://doi.org/10.17503/agrivita.v40i0.469>[Accessed 4 September 2018].
8. Institut Pertanian Bogor (IPB). 2005. *Pasar Domestik dan Ekspor Produk Tanaman Obat*. IPB Bogor: Bogor Agricultural University.
9. Singh, S., Gautam, A., Sharma, A. and Batra, A. 2010. *Centella asiatica* (L.): A plant with immense medicinal potential but threatened. *Int. J. Pharm. Sci. Res.*,4(2):9-17. Available from: <http://globalresearchonline.net/journalcontents/volume4issue2/Article%20003.pdf> [Accessed 8 September 2018].
10. Alqahtani, A., Tongkao-on, W., Li, K.M., Razmovski-Naumovski, V., Chan, K. and Li, G.Q. 2015. Seasonal variation of triterpenes and phenolic compounds in Australian *Centella asiatica* (L.) Urb. *Phytochem. Anal.*, 26(6):436-443. Available from: <https://doi.org/10.1002/pca.2578> [Accessed 20 December 2018].
11. Kim, O.T., Kim, M.Y., Hong, M.H., Ahn, J.C. and Hwang, B. 2004. Stimulation of asiaticoside accumulation in the whole plant cultures of *Centella asiatica* (L.) Urban by elicitors. *Plant Cell Rep.*, 23(5): 339-344. Available from: <https://doi.org/10.1007/s00299-004-0826-7> [Accessed 2 September 2018].
12. Lambert, E., Faizal, A. and Geelen, D. 2011. Modulation of triterpene saponin production: in vitro cultures, elicitation, and metabolic engineering. *Appl. Biochem. Biotechnol.*, 164(2):220-237. Available from: <https://doi.org/10.1007/s12010-010-9129-3> [Accessed 11 September 2018].
13. Bonfill, M., Mangas, S., Moyano, E., Cusido, R.M. and Palazón, J. 2011. Production of centellosides and phytosterols in cell suspension cultures of *Centella asiatica*. *Plant Cell Tissue Organ Cult.*,104(1):61-67. Available from: <https://doi.org/10.1007/s11240-010-9804-7> [Accessed 1 January 2019].
14. Mangas, S., Moyano, E., Osuna, L., Cusido, R.M., Bonfill, M. and Palazón, J. 2008. Triterpenoid saponin content and the expression level of some related genes in calli of *Centella asiatica*. *Biotechnol. Lett.*, 30(10):1853. Available from: <https://doi.org/10.1007/s10529-008-9766-6> [Accessed 1 September 2018].
15. Ghulamahdi, M., Aziz, S.A. and Bermawie, N. 2007. *Evaluasi Karakter Morfologi, Fisiologi dan Genetik Pegagan Mendukung Standarisasi Mutu Pegagan*. Bogor, Indonesia. Bogor Agricultural University.
16. Gupta, A., Verma, S., Kushwaha, P., Srivastava, S. and Aks, R. 2014. Quantitative estimation of asiatic acid, asiaticoside & madecassoside in two accessions of *Centella asiatica* (L) Urban for morpho-chemotypic variation. *Indian J. Pharm. Educ.*48:75-79. Available from: <https://www.ijper.org/sites/default/files/10.5530ijper.48.3.9.pdf> [Accessed 25 February 2019].
17. Vinolina, N.S. and Siregar, L.A. 2012. Morphology of leaves and content of secondary metabolites asiaticoside in some accession of pegagan (*Centella asiatica* L. Urban) in North Sumatera. *Proceedings of The Annual International Conference, Syiah Kuala University-Life Sciences & Engineering Chapter.*, 2(1): 200-205. Available from: <http://www.jurnal.unsyiah.ac.id/AICS-SciEng/article/view/1768/1665> [Accessed 12 December 2018].
18. Jain, P.K. and Agrawal, R.K. 2008. High performance liquid chromatographic analysis of asiaticoside in *Centella asiatica* (L.) Urban. *Chiang Mai J. Sci.*35(3): 521-525. Available from: http://it.science.cmu.ac.th/ejournal/journalDetail.php?journal_id=278 [Accessed 20 December 2018].
19. Mangas, S., Bonfill, M., Osuna, L., Moyano, E., Tortoriello, J., Cusido, R.M., Piñol, M.T. and Palazón, J. 2006. The effect of methyl jasmonate on triterpene and sterol metabolisms of *Centella*

- asiatica, *Ruscus aculeatus* and *Galphimia glauca* cultured plants. *Phytochemistry*.67(18):2041-2049. Available from: <https://doi.org/10.1016/j.phytochem.2006.06.025> [Accessed 1 September 2018].
- 20.Jamil, S.S., Nizami, Q. and Salam, M. 2007. *Centella asiatica* (Linn.) Urban—a review. *Natural Product Radiance*, 6(2):158-170. Available from: <http://hdl.handle.net/123456789/7855> [Accessed 26 February 2019].
- 21.Zainol, M.K., Abd-Hamid, A., Yusof, S. and Muse, R. 2003. Antioxidative activity and total phenolic compounds of leaf, root and petiole of four accessions of *Centella asiatica* (L.) Urban. *Food Chemistry*.81(4):575-581. Available from: [https://doi.org/10.1016/S0308-8146\(02\)00498-3](https://doi.org/10.1016/S0308-8146(02)00498-3) [Accessed 29 December 2018]
- 22.Rosenthal, G.A. and Berenbaum, M.R. 2012. *Herbivores: their interactions with secondary plant metabolites: ecological and evolutionary processes*. Vol 2: Academic Press.
- 23.Bennett, R.N. and Wallsgrave, R.M. 1994. Secondary metabolites in plant defence mechanisms. *New phytologist*.127(4):617-633. Available from: <https://doi.org/10.1111/j.1469-8137.1994.tb02968.x> [Accessed 29 December 2018].Page Break

FIGURES

FIGURES




Figure 1. Asiaticoside concentration ($\mu\text{g/mL}$) in leaves (KAD) and roots (KAA) of Pegagan harvested at 56 (U_1), 70 (U_2) dan 84 (U_3) DAP. Significant differences are indicated by different letters above the bars (F-tests, $p < 0.05$).




Figure 2. Madecassoside concentration ($\mu\text{g/mL}$) in leaves (KMD) and roots (KMA) of Pegagan harvested at 56 (U_1), 70 (U_2) dan 84 (U_3) DAP. Significant differences are indicated by different letters above the bars (F-tests, $p < 0.05$).




Figure 3. Asiatic acid concentration ($\mu\text{g/mL}$) in leaves (KAAD) and roots (KAAA) of Pegagan harvested at 56 (U_1), 70 (U_2) dan 84 (U_3) DAP. Significant differences are indicated by different letters above the bars (F-tests, $p < 0.05$)

TABLE

Table 1. Wet weight to dry weight of leaves and roots of Pegagan harvested at 56 (U_1), 70 (U_2) dan 84 (U_3) DAP

Treatment	WWL (g)	DWL (g)	WWR (g)	DWR (g)
U1	77.623 c	13.665 c	66.158 c	12.291 c
U2	191.206 b	17.149 b	188.894 b	22.642 b
U3	201.211 a	26.206 a	200.733 a	31.553 a

Note: Different values in the same column indicated significant difference at $p < 0.05$. WW(L/R) = wet weight of leaves/roots; DW(L/R) = dry weight of leaves/roots

AUTHOR CONTRIBUTION

Name of the author and e-mail ID	Types of contribution
Noverita Sprinse Vinolina noveritasitumorang@yahoo.com	Primary author, conducting the overall research arrangement and cultivation of <i>Centella Asiatica</i> including grant proposal writing and submission, seed planting, supervision of field preparation, harvest, UFLC analysis of secondary metabolite components. Writing the manuscript and editing contributions from co-authors

Luthfi Siregar luthfi2004@yahoo.com	Co-author, carrying out the collection of mother plants, preparation of samples for UFLC analysis and contributed in writing the manuscript.
Biliter Sirait dapejel@yahoo.com	Co-author, carrying out the collection of mother plants, supervision of field preparation, preparation of samples for UFLC analysis and contributed in writing the manuscript.
Nurhayati nurhayati@uisu.ac.id	Co-author, carrying out the collection of mother plants, supervision of field preparation, preparation of samples for UFLC analysis and contributed in writing the manuscript.
Enny Adelina ennyadelina@gmail.com	Co-author, analysis data dan contributed in writing the manuscript.
Laila Nazirah laila_nazirah@yahoo.co.id	Co-author, analysis data dan contributed in writing the manuscript.