

THE 5th ANNUAL INTERNATIONAL CONFERENCE SYIAH KUALA UNIVERSITY
(AIC - UNSYIAH)
IN CONJUNCTION WITH
THE 8th CHEMICAL ENGINEERING ON SCIENCE AND APPLICATION
(ChESA)

LIFE SCIENCES CHAPTER



AAC Dayan Dawood, Darussalam - Banda Aceh, Indonesia
September 9 - 11, 2015



**Proceeding of the 5th Annual International Conference (AIC) Syiah
Kuala University 2015 in Conjunction with the 8th Chemical
Engineering Science & Applications (ChESA)
9-11 September 2015, Banda Aceh - Indonesia**

Editor in Chief : Dr. drh. Ummu Balqis, M.Si.
(Syiah Kuala University, Indonesia)

Editors :

- Prof. Dr. drh. Darmawi, M.Si.**
(Syiah Kuala University, Indonesia)
- Prof. Dr. Ir. Samadi, M.Sc.**
(Syiah Kuala University, Indonesia)
- Prof. Roselyn M. Butalid**
(Mindanao State University, Philippines)
- Associate Prof. Dr. Ororat Mongkolporn, Ph.D**
(Kasetsart University, Thailand)
- Prof. Dr. Dr. h.c. Evamarie Hey-Hawkins**
(Universität Leipzig, Germany)
- Prof. Dr. med. vet. Jean-Michel Hatt, MSc, Dipl ACZM, ECZM (Avian)**
(University of Zurich, Switzerland)
- Dr. Kazuaki Syutsubo**
(National Institute for Environmental Studies, Japan)
- Prof. Dr. Ir. H. Hasanuddin, M.S.**
(Syiah Kuala University, Indonesia)
- drh. Al Azhar, M. Kes.**
(Syiah Kuala University, Indonesia)

ISSN: 2089-208X
Copyright © 2015
Printed September 2015

CONTENT

EDITORIAL BOARD		Pages
MESSAGE FROM THE RECTOR		
MESSAGE FROM THE CHAIRMAN		
No	Scientific Paper	
	Keynote and Invited Speaker	1
1	SNP Mapping to Locate Anthracnose Resistance in <i>Capsicum</i> spp. <i>Orarat Mongkolporm (Thailand)</i>	2-5
2	Development of simple and cost-effective treatment system for municipal wastewater <i>Takashi Onodera, Kazuaki Syutsubo (Japan)</i>	6
3	The Importance of Chemistry for Nanotechnology <i>Evamarie Hey-Hawkins (Germany)</i>	7
4	Hybrid Catalytic-Plasma Reactor for Energy and Chemical Conversions <i>I. Istadi (Indonesia)</i>	8
5	Photocatalytic oxidation in phenol removal using Ru/TiO ₂ and Ru/Al ₂ O ₃ catalysts <i>Syaifulah Muhammad (Indonesia)</i>	9
	Agriculture	10
1	Microstructure and Optimization of Sweet Potato-Rice Simulated Chips <i>Adil Basuki Ahza (Indonesia), M. Arief Munandar(Indonesia), and Subarna Suryatman (Indonesia)</i>	11-15
2	Some Rice Varieties Tolerant to Nitrogen Deprivation <i>Muhammad Hatta (Indonesia), Rosmayati (Indonesia), Chairani Hanum (Indonesia), and Hamidah Hanum (Indonesia)</i>	16-19
3	Effect of Leaf Extract Sirih (<i>Piper betle</i> L.) and on Future Soaking Time Freshness of Flowers Rose (<i>Rosa sinensis</i> L.) <i>Devri Wahyuni (Indonesia), Rita Hayati (Indonesia), and Nurhayati (Indonesia)</i>	20-24
4	How to restore the Tropical Peat Swamp Forest in Aceh Province, Indonesia <i>Hairul Basri (Indonesia), Sufardi (Indonesia), Suglanto (Indonesia), Syamaun A Ali (Indonesia), Khalrullah (Indonesia), and Ahmad Reza Kasuri (Indonesia)</i>	25-28
5	Effect of Corn Extract and Solution of PEG to Increase Viability and Vigour of Sweet Corn Seed Deterioration at Different Incubation Periods <i>Herme Julendri (Indonesia), Hafimuzayudah (Indonesia), and Hesenuddin (Indonesia)</i>	29-32
6	Fig Pollinating Wasp Transfers Nematodes into Figs of <i>Ficus racemosa</i> in Sumatra, Indonesia <i>Jauharlina (Indonesia), Rina Sriwati (Indonesia), Yusmaini (Indonesia), Natsumi Kanzaki (Japan), Stephen Compton (South Africa)</i>	33-37
7	Physiological Seed Performance of Local Aceh and National Release Variety of Rice (<i>Oryza sativa</i> L.) to Water Stress <i>Cut Nur Ichsana (Indonesia), Bakhtiar (Indonesia), Efendi (Indonesia), and Sabarudin (Indonesia)</i>	38-42
8	Application of Soil Conservation in Oil Palm Plantation <i>Halus Satriawan (Indonesia), Zahrul Fuady (Indonesia), and Zara Yunizar (Indonesia)</i>	43-46
9	Genetic Characterization and Estimation Variety Eight of Papaya Genotype <i>Siti Hafsa (Indonesia), Yuznizar (Indonesia), and Firdaus (Indonesia)</i>	47-49
10	Adaptation of Some Clones of Sweet Potato (<i>Ipomoea batatas</i> L.) in the Lowlands and the Use of Manure <i>Mardhiah Hayati (Indonesia), Sabaruddin (Indonesia), Efendi (Indonesia), Ashabul Anhar (Indonesia), Ely Kesumawati (Indonesia), and NurKamalia (Indonesia)</i>	50-54
11	Investigation of the Effect of Heat Moisture Treatment on Local Sweet Potato Starch Characteristics <i>Zeldiyah (Indonesia), and Lukmanul Hakim (Indonesia)</i>	55-57
12	Rejuvenation and Characterization of Local Rice Germplasm (<i>Oryza sativa</i> L.) Under Organic Cultivation System <i>Efendi (Indonesia), Bakhtiar (Indonesia), Sabaruddin Zakaria (Indonesia), and Syafruddin (Indonesia)</i>	58-62
13	Roots Bioassay of Upland Rice Varieties on Several Soil Moisture Gradients <i>Syarifa Mayly (Indonesia), A. Rauf (Indonesia), Chairani Hanum (Indonesia), and Hamidah Hanum (Indonesia)</i>	63-66
14	Compost Oil Palm and Indigenous Endophytic Fungi Effect on Basal Stem Rot in Oil Palm Seedling <i>Lisnawita (Indonesia), Hamidah Hanum (Indonesia), and Ahmad Rafiqi Tantawi (Indonesia)</i>	67-70
15	Colonization Ability of Biological Control Agent <i>Trichoderma</i> spp on <i>Cucur</i> Poul and Seedling <i>Rina Sriwati (Indonesia), Tjut Khamzurni (Indonesia), Elvira Iskandar (Indonesia), and</i>	71-75

Compost Oil Palm and Indigenous Endophytic Fungi Effect on Basal Stem Rot in Oil Palm Seedling

¹Lisnawita, ²Hamidah Hanum, and ³Ahmad Rafiqi Tantawi

^{1,2}Program Study of Agroecotechnology, Faculty of Agriculture, University of Sumatera Utara, Padang Bulan, Medan 20155, Indonesia;

³Program Study of Agrotechnology, Faculty of Agriculture, University of Medan Area, Medan 20223, Indonesia

*Corresponding Author: itamuis@yahoo.com

Abstract

Basal stem rot (BSR) caused by *Ganoderma* is an important disease of oil palm in North Sumatra, Indonesia. Control of BSR is complex, because the disease occurs in the dynamic environment at the interface of the roots with soil. Compost with endophytic fungi as a biological fertilizer offer possible advantages and eventually suppressed *Ganoderma* growth. A study using two compost oil palm (empty fruit bunch and palm oil midrib-leaf compost) containing indigenous endophytic fungi (*Trichoderma koningii*, *T. viride*, and *Aspergillus* sp.) singly and in a mixture was conducted to determine potential of them to control *Ganoderma*. The result showed that 16 weeks after artificial inoculation, all treatments aren't significant difference reduced BSR incidence but compost containing endophytic fungi have potential to inhibit the pathogen showed BSR incidence in a mixture treated lower than singly treated seedlings.

Key words: *Aspergillus* sp., compost, *Ganoderma*, oil palm, *Trichoderma koningii*, *T. viride*

Introduction

Oil palm (*Elaeis guineensis*) is one of the largest plantation commodity industries in Indonesia. Currently, Indonesia is one of the world's largest producer and exporter of oil palm. Areas of oil palm have increased from 135.000 hectares in 1970 to 8.91 million in 2011 (Pusat Data dan Sistem Informasi Pertanian, 2013), which producing an average palm oil yield approximately 31 tonnes in 2014 (<http://www.indonesia-investments.com>, 2015). One of province with the second largest oil palm planted area in Indonesia is North Sumatra with oil palm planted area is 1.2 million hectares (Direktorat Jenderal Perkebunan, 2012).

One major problem in oil palm is basal stem rot (BSR) disease, caused by *Ganoderma* spp., which is a serious threat to the oil palm industry in South East Asian countries, especially Malaysia and Indonesia. The disease is lethal and in the past few decades has been spreading rapidly. The disease spreads through root to root contact. During planting any little piece of rotting trunk is removed as this could be a potential source of *Ganoderma* in the current stand (Hasar and Turner, 1998). The disease progresses slowly, but eventually all infected plants become dead. This is because the disease cannot be detected at the early stages.

Field control of BSR by contact chemicals have not been very successful (Scepena et al., 2000) even with those *in vitro* screened effective against the fungus (Khairudin, 1990; Teh, 1996). The current interest in reducing the application of chemical fertilizer and increasing demand for combined effects of beneficial fungi and organic compost can have great impact on crop production and sustainable agriculture. Compost with endophytic fungi as a biological fertilizer offer possible advantages and eventually suppressed *Ganoderma* growth. A study using two compost oil palm (empty fruit bunch and palm oil midrib-leaf compost) containing indigenous endophytic fungi (*Trichoderma koningii*, *T. viride*, and *Aspergillus* sp.) singly and in a mixture was conducted to determine potential of them to control *Ganoderma*.

Materials and Methods

Preparation of *Ganoderma* inoculum on rubber wood blocks

Fresh rubber wood blocks (12 cm x 6 cm x 6 cm, weighing approximately 450 g) were washed with water and autoclaved for 20 min at 121 °C. One block each was put in heat-resistant polypropylene bags (15 cm x 33 cm x 0.05 mm thick material) and 100 mL of potato dextrose agar (PDA) added as supplementary nutrient for *Ganoderma*. Since inoculation of the blocks with *Ganoderma* plugs can introduce contamination, the bags were closed by drawing its open end through a 4 cm diameter polyvinyl chloride (PVC) tubing 2 cm long and the remaining hole plugged with cotton. The bags with rubber wood blocks and PDA were autoclaved at 121 °C for 30 min. After sterilization and cooling, the

rubber wood block in the bag was rotated to ensure that it was well covered with the agar before the latter solidified. When the agar had solidified, five pieces five day-old *Ganoderma* culture taken with cork borer and inoculated on each rubber wood block. The inoculated blocks in the bags were then incubated in a dark cabinet at $28 \pm 2^\circ\text{C}$ for three weeks until fully colonized by *Ganoderma* mycelium.

Preparation of endophytic fungi

Trichoderma koningii, *Trichoderma viride*, and *Aspergillus* sp. were isolated from the oil palm roots in Gunung Melayu, Asahan, North Sumatra and cultured on PDA medium. All of endophyt fungi were prepared using rice medium, then formulated with zeolite powder (1:9 /v:v).

Application of compost oil palm containing indigenous endophytic fungi

Experiments carried out in the glasshouse at Faculty of Agriculture, University of Sumatra Utara, Medan, Indonesia, using a factorial randomized block design. A total of 72 healthy oil palm seedlings three months old were used in this experiment. 300 g of compost single or combination with 30 g of endophytic fungi inoculated before or after artificial inoculation of *Ganoderma* on oil palm seedlings.

Effect of compost oil palm and indigenous endophytic fungi on BSR incidence

At the end of the experiment (4 months/16 weeks), the oil palm seedlings were split longitudinally to observe for root and stem decay and to visually assess the severity of the internal symptoms based on the proportion of root and bole tissues damaged by *Ganoderma*. The estimation was based on the following modified scale (Abdullah *et al.*, 2003; Ilias 2000). 0 = Healthy plants, 1 = Appearance of white fungal mass on any part of plants, with or without chlorotic leaves, 2 = Appearance of basidioma on any part of plants with chlorotic leaves (1-3 leaves), 3 = Formation of basidioma on any part of plants with chlorotic leaves (> 3 leaves), 4 = Formation of well-developed basidioma and the plants dried. Disease severity (DS) for the internal symptoms was calculated by the formula of Liu *et al.* (1995) as under. $DS_{(internal)} = (\text{number of seedlings in the rating} \times \text{rating number}) / (\text{total number of seedlings assessed} \times \text{highest rating}) \times 100$. The disease incidence and disease severity percentages were transformed by arcsine and analyzed using ANOVA with the means compared by the LSD ($P \leq 0.05$).

Results and Discussion

Artificial infection of plants by contact with the inoculum block carrying *Ganoderma* is an effective strategy for inducing the infection. The assessment of disease severity was based on the sequential progression of disease using index values 0-4. Lower disease severity indicates the effectiveness of the treatments in suppressing *Ganoderma* BSR in oil palm through *in vivo* assessment. In general, 16 weeks after artificial inoculation, all treatments aren't significant difference reduced BSR incidence but compost containing endophytic fungi have potential to inhibit the pathogen showed BSR incidence in a mixture treated lower than singly treated seedlings (Table 1). This study showed that compost containing endophytic fungus (*T. koningii*, *T. viride*, and *Aspergillus* sp.) have a good potential as biological agents to inhibit *Ganoderma*.

Table.1. Disease severity of oil palm plants at BSR 16 weeks after artificial inoculation

Treatments	Disease severity (%)
K0W1E0	33.3
K0W1E1	33.3
K0W1E2	25.0
K0W1E3	25.0
K0W2E0	25.0
K0W2E1	25.0
K0W2E2	25.0
K0W2E3	25.0
K1W1E0	25.0
K1W1E1	8.3
K1W1E2	16.7
K1W1E3	16.7
K1W2E0	25.0
K1W2E1	16.7
K1W2E2	16.7

K1W2E3	8.3
K2W1E0	25.0
K2W1E1	16.7
K2W1E2	16.7
K2W1E3	8.3
K2W2E0	25.0
K2W2E1	16.7
K2W2E2	8.3
K2W2E3	16.7

Note : K0 : without compost,
 K1 : empty fruit bunch compost,
 K2 : palm oil midrib-leaf compost,
 W1 : inoculation *Ganoderma* before endophytic fungi
 W2 : inoculation *Ganoderma* after endophytic fungi
 E0 : without endophytic fungi
 E1 : *T. viride*
 E2 : *T. koningii*
 E3 : *Aspergillus* sp.

The composition of the microorganisms in composts is affected by the chemistry of the materials from which the compost is prepared (Castaño *et al.*, 2011). Composts with high lignocellulosic substances (tree barks) are mostly colonized by *Trichoderma* spp. In contrast, grape pomace, with low cellulosic substances and high sugars, becomes colonized by *Penicillium* spp. and *Aspergillus* spp. (Kutter *et al.* 1983; Gorodecki and Hadar 1990). In composts, in addition to the well-documented effects of microbial populations against several soil-borne diseases, induction of resistance has also been reported as an additional biocontrol mechanism against both foliar and root diseases (Zhang *et al.* 1996, 1998; Kavroulakis *et al.* 2005; Ntougias *et al.*, 2008). Reduction in disease severity was demonstrated against *Septoria lycopersici* in tomato (Kavroulakis *et al.*, 2005), bacterial leaf spot in radish caused by *Xanthomonas campestris* pv. *Armoraciae* (Krause *et al.*, 2003), anthracnose in cucumber caused by *Colletotrichum orbiculare* (Zhang *et al.*, 1996) and *Botrytis cinerea* in cucumber and melon (Segarra *et al.* 2007; Yogeve *et al.*, 2010). Moreover compost water extract root treatment induce systemic resistance to anthracnose caused by *C. orbiculare* in cucumber and *C. coccodes* in pepper (Sang and Kim 2011).



Figure 1. Signs and symptoms of BSR, caused by *Ganoderma*; A, B. necrotic lesion on infected plant (arrow), C. The leaves of infected plants were chlorotic (a), and healthy leaves (b)

The ideal of biocontrol strategy attempts to introduce or promote the activity of biocontrol agents only when and where they are needed or are most effective, and minimises wasteful application of inoculum on non-target habitats. Different mechanisms have been suggested as being responsible for the effects of biocontrol agents; they include competition for space and nutrients, secretion of chitinolytic enzymes, mycoparasitism and production of an inhibitory compound (Haram *et al.*, 1996; Zimand *et al.*, 1996). Thus, the timing of when to apply the treatments is important due to the

probability of changes brought about by the biocontrol agent, e.g. by the latter's dominating requirements for space and nutrients.

Artificial inoculation and following very close contact with the root surface, penetration of the root epidermis and exodermis occurred. Heavily infected tissue within the root was evident on sectioning by brown discoloration primarily in the cortex, but in some instances *Ganoderma* mycelium emerged through ruptures in the epidermis of infected roots. The cross-section of an infected plant stem showed a necrotic lesion at the vascular region (Figure 1A and B). Immediately in advance of infection was often a small area of yellow tissue which has been observed by other authors including Darmono (1998). The leaves of infected plants were chlorotic and had a white fungal mass on any part of the plants, then formed basidioma before the plants dried (Figure 1C).

Conclusion

Amount of 16 weeks after artificial inoculation, all treatments aren't significant difference reduced BSR incidence but compost (empty fruit bunch and palm oil midrib-leaf compost) containing indigenous endophytic fungi (*Trichoderma koningii*, *T. viride*, and *Aspergillus* sp.) have potential to inhibit the pathogen showed BSR incidence in a mixture treated lower than singly treated seedlings.

Acknowledgement

This research was supported by KEMENRISTEK DIKTI through The National Competitive Grants Scheme MP3EI (2014-2015). Many thanks for the financial support.

References

- Abdullah F., Ilias G.N.M., Nelson M., Nur Ain Izzati M.Z., Umi Kalsom Y. (2003): Disease assessment and the efficacy of *Trichoderma* as a biocontrol agent of basal stem rot of oil palms. *Research Bulletin Science Putra*, 11: 31-33.
- Castaño R., Borrero C., Avilés M. (2011). Organic matter fractions by SP-MAS ¹³C NMR and microbial communities involved in the suppression of *Fusarium* wilt in organic growth media. *Biological Control* 58, 286-293
- Direktorat Jenderal Perkebunan. 2012. Luas areal kelapa sawit menurut Provinsi di Indonesia, 2008-2012. www.deptan.go.id. (Diunduh 13 Desember 2013).
- Darmono TW, 2000. *Ganoderma* in oil palm in Indonesia: Current status and prospective use of antibodies for the detection of infection. In: (J. Flood, P. Bridge & M. Holderness, eds) *Ganoderma Diseases of Perennial Crops*: Wallingford: CABI Publishing. 249-66.
- Gorodecki B., Hader Y. (1990). Suppression of *Rhizoctonia solani* and *Sclerotium rolfsii* in container media containing composted separated cattle manure and composted grape marc. *Crop Protection* 9, 271-274
- <http://www.indonesia-investments.com>, 2015. **Food and Agriculture Organization of the United Nations, Indonesian Palm Oil Producers Association (Gapki) and Indonesian Ministry of Agriculture**. (Diunduh 20 Agustus 2015).
- Hasan Y., Turner P.D. (1998). The comparative importance of different oil palms tissues as infection sources for basal stem rot in replanting. *The Planter*. 74:119-135.
- Ilias G.N.M. (2000). *Trichoderma* and its efficacy as a bio-control agent of basal stem rot of oil palm (*Elaeis guineensis* Jacq.). [Ph.D. Thesis.] Universiti Putra Malaysia, Selangor, Malaysia.
- Khairudin H. (1990). Basal stem rot of oil palm: incidence, etiology and control. MSc thesis. Universiti Pertanian Malaysia, Selangor, Malaysia.
- Kuter G.A., Nelson E.B., Holtink H.A.J., Madden L.V. (1983). Fungal populations in container media amended with composted hardwood bark suppressive and conducive to *Rhizoctonia damping-off*. *Phytopathology* 73, 1450-1456.
- Kavroulakis N., Ehallotis C., Ntougias S., Zervakis G.I., Papadopoulou K.K. (2005). Local and systemic resistance against fungal pathogens of tomato plants elicited by a compost derived from agricultural residues. *Physiological and Molecular Plant Pathology* 66, 163-174.
- Krause M.S., De Ceuster T.J.J., Tiquia S.M., Michel F.C.J.R., Madden L.V., Hol-tink H.A.J. (2003). Isolation and characterization of rhizobacteria from com-posts that suppress the severity of bacterial leaf spot of radish. *Phytopathology* 93, 1116-1123.
- Ntougias S., Papadopoulou K.K., Zervakis G.I., Kavroulakis N., Ehallotis C. (2008). Suppression of soil-borne pathogens of tomato by composts derived from agro-industrial wastes abundant in Mediterranean regions. *Biology and Fertility of Soils* 44, 1081-1090.
- Pusat Data dan Sistem Informasi Pertanian, 2013. Kelapa Sawit. Informasi Ringkas Komoditas Perkebunan. www.deptan.go.id. (Diunduh 13 Desember 2013).
- Soepena, H., R.Y. Purba and S. Pawirosukarto, 2000. A control strategy for basal stem rot (*Ganoderma*) on oil palm. In: Flood, J. et al. (eds.), *Ganoderma Disease of Perennial Crops*, p: 83-8. CAB International, UK.
- Sang M.K., Kim K.D. (2011). Biocontrol activity and primed systemic resistance by compost water extracts against anthracnoses of pepper and cucumber *Phytopathology* 101, 732-740.
- Segarra G., Casanova E., Borrero C., Avilés M., Trillas I. (2007). The suppressive effects of composts used as growth media against *Botrytis cinerea* in cucumber plants. *European Journal of Plant Pathology* 117, 393-402.