

COMPARISON IMMUNOHISTOCHEMISTRY EXPRESSION OF DESIDUAL NATURAL KILLER (dNK) IN SEVERE PREECLAMPSIAAND NORMAL PREGNANCY

by Muara Panusunan Lubis

Submission date: 09-Jul-2020 10:59AM (UTC+0700)

Submission ID: 1355265105

File name: ATURAL_KILLER_dNK_IN_SEVERE_PREECLAMPSIAAND_NORMAL_PREGNANCY.pdf (243.53K)

Word count: 1978

Character count: 10313

COMPARISON IMMUNOHISTOCHEMISTRY EXPRESSION OF DESIDUAL NATURAL KILLER (dNK) IN SEVERE PREECLAMPSIA AND NORMAL PREGNANCY

MUARA P. LUBIS¹

¹Fetomaternal Division, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Sumatera Utara
Email: mploebeis@yahoo.com

Received: 25 Jan 2020, Revised and Accepted: 18 Mar 2020

ABSTRACT

Objective: Preeclampsia characterized systematically by extensive vascular endothelial dysfunction and microangiopathy on mother, dNK is very important for the success of placentation. They are the key mediator of maternal immune system interactions with fetal cells. dNK cells are also involved in the modulation of EVT and the remodeling of spiral arteries.

Methods: Analytic research with cross-sectional study, with samples of pregnant women who suffer from severe PE and term pregnancy which came to H. Adam Malik Hospital and Networking Hospital, November 2015-April 2016. The samples are 46 women, who met the inclusion criteria.

Results: Immunohistochemistry examination dNK cell in the severe PE case group and control group, statistically found $p < 0,05$. dNK placenta expression in the severe preeclampsia case group gives an overview of expression with a mean of 2.55 ± 2.31 , while the control group of normal pregnancy had higher mean is 8.66 ± 3.16 .

Conclusion: The examination of immunohistochemistry of dNK cells showed there is a significant difference in the expression of Immunohistochemistry dNK cells between severe PE case group and non severe PE.

Keywords: Severe preeclampsia, Normal pregnancy, Expression of dNK cell

© 2020 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)
DOI: <http://dx.doi.org/10.22159/ijcpr.2020v12i3.38306> Journal homepage: <https://innovareacademics.in/journals/index.php/ijcpr>

INTRODUCTION

Preeclampsia is systemically characterized by widespread vascular endothelial dysfunction and microangiopathy in the mother, but not in the fetus. At present it is believed that preeclampsia starts with abnormalities in the development of blood vessels in the placenta that cause an effect on the maternal endothelium [1, 4, 5].

Knowledge of cellular and molecular processes of human trophoblast invasion is based on *in vitro* research and animal models; there is evidence that decidual natural killer (dNK) cells are very important in successful placentation. They are key mediators of the interaction of the mother's immune system with fetal cells. dNK cells are also involved in modulation of invasion of extravillous trophoblasts (EVT) and remodeling of the maternal spiral arteries. They express various surface receptors and signaling molecules, including cytokines, chemokines, and growth factors, and their function in modulating EVT migration, invasion, and change from epithelial phenotype to endothelial are beginning to be revealed [6].

MATERIALS AND METHODS

This study is a comparative analytic study with a cross-sectional study design conducted at General Hospital. H. Adam Malik Medan,

RSUD dr. Pirngadi Medan, and Faculty of Medicine, University of North Sumatra Networking Hospital from November 2015 to April 2016. The sample of the study was 23 pregnant women suffering from PEB and normal pregnant women with a term gestational age for each group who met the inclusion criteria, namely pregnant women aged 18-35 y, singleton pregnancy, severe preeclampsia, term and exclusion criteria ie damaged placenta samples; who came for pregnancy control to General Hospital H. Adam Malik Medan, RSUD dr. Pirngadi Medan, Faculty of Medicine, University of North Sumatra Networking Hospital.

Work arrangement

Taken in the middle of the maternal side opposite the insertion site of the umbilical cord on the fetal surface with a size of 1.5 x 1.5 x 1 cm. Then fixed in 10% formalin liquid with a time limit of 24-48 h, dried with ethanol and cleaned with xylol and implanted in a paraffin block. Paraffin is cut 7µm. Stained with Hematoxylin and Eosin (H and E). Immunohistochemically stained for CD56 to detect dNK cells using CD56 anti antibodies. The preparations were interpreted by two Anatomical Pathologists. The results of the preparation of these preparations are carried out statistical analysis with Statistical Product and Service Solutions (SPSS) Ver. 18

RESULTS

Characteristics of research subjects

Table 1: Distribution characteristics of research subjects in severe preeclampsia case group and normal pregnancy group

Characteristics	Research subjects		Normal pregnancy	%	Total	%	p value
	Severe preeclampsia	%					
Age (years old)							
15-25	3	13	8	34,8	11	23,9	0,062**
26-35	16	69,6	15	65,2	31	67,4	
>35	4	17,4	0	0	4	8,7	
Parity							
Primigravida	7	30,4	5	21,7	12	26,1	0,611*

Secundigravida	6	26,1	9	39,1	15	32,6	
Multigravida	10	43,5	9	39,1	19	41,3	
Birth Weigth							
Low Birth Weight	5	21,7	0	,0	5	10,9	0,058***
Normal	18	78,3	23	100	41	89,1	
APGAR Score							
Good	23	100	23	100	46	100	
PROTEINURIA							
(+2)	9	39,1	0	0	9	19,6	
(+3)	11	47,8	0	,0	11	23,9	
(+4)	3	13,0	0	,0	3	6,5	0,001**
Negative	0	0	23	100	23	50	
Total	23	100	23	100	46	100	

*Chi-square test **Fisher Exact test ***Continuity Correction

Table 2: Differences in dNK cell expression between severe preeclampsia case group and normal pregnancy group

Research subjects	N	CD56 expressions		p value
		Mean	Std. deviasi	
Severe Preeclampsia	23	2,55	2,31	0,031
Normal Pregnancy	23	8,66	3,16	

Table 3: Differences in dNK cell expression between severe preeclampsia case group and normal pregnancy group based on Normal birth weigth and low birth weight (LBW)

Birth weight	N	CD56 expressions		p value
		Mean	Std. deviasi	
Normal	41	6,9451	5,20322	0,003
LBW	5	2,4000	2,00468	

Table 4: Differences in expression of dNK cells based on proteinuria

Research subjects	N	CD56 expressions		p value
		Mean	Std. deviasi	
Proteinuria				
Negative	23	9,6957	4,24997	0,001
(+2)	9	3,8889	4,90075	
(+3)	11	2,7045	2,93645	
(+4)	3	3,0000	3,46410	




Fig. 1: Boxplot showing mean number of dNK cells in placenta with severe preeclampsia and normal pregnancy

DISCUSSION

Immunohistochemistry examination results

The results of this study found that the results of immunohistochemistry dNK cells examination in the severe preeclampsia case group generally gave a description of expression with a mean of 2.55 ± 2.31 while in the normal pregnancy control group had a higher mean with 8.66 ± 3.16 . Statistically obtained p value < 0.05 which indicates there is a significant difference in the expression of immunohistochemistry dNK cells between severe preeclampsia case group and normal pregnancy group.

In another study by Charles *et al.* stated that dNK cells in women with preeclampsia were significantly less than controls (normotension). Closely related between dNK cells and blood vessels, dNK cells trigger angiogenic factors on the effects of IFN- γ to facilitate remodeling of normal blood vessels [57, 70, 71]. Rieger *et al.* in their study found that there was a significant relationship between CD56+/CD16+dNK cell counts that increased in the normal pregnancy group than in severe preeclampsia group (7.3% vs 5.3%, $p < 0.02$) [58]. Williams *et al.* observed that CD56+NK cells ($p = 0.01$) decreased in placental bearing biopsies in PE pregnancy women compared to normal third trimester pregnancy [59].

CONCLUSION

Immunohistochemistry examination of dNK cell in severe preeclampsia case group generally gave a picture of expression with a mean of 2.55 ± 2.31 while in the normal pregnancy control group had a higher mean with 8.66 ± 3.16 . Which showed significant differences in the expression of immunohistochemistry dNK cell between severe preeclampsia case group and normal pregnancy group.

ACKNOWLEDGEMENT

On this special joyful occasion let me express my deepest gratitude to: Dr. dr. Makmur Sitepu, M. Ked (OG), SpOG. K, Prof. dr. Daulat Sibuea, SpOG. K, dr. Makmur Sitepu, M. Ked (OG), SpOG. K, dr. Jessy Chrestella, M. Ked (PA), SpPA, dr. Surya Darma who has helped a lot in this research. Beloved wife, Dr. Fithria Aldy, M. Ed (Oph), Sp. M, and our three children Gandisyah Khalisa Mahira Lubis, Gandira Alisha Hanifa Lubis, and M. Chairuddin Martua Lubis. May Allah Subhanahu wa ta'ala bestow His mercy and guidance on us all. In addition, researchers are aware of the need for further research so that in the future it can become a reference basis for immunological therapy in severe preeclampsia

FUNDING

Nil

AUTHORS CONTRIBUTIONS

All the authors have contributed equally.

CONFLICT OF INTERESTS

There is no conflict of interest in this research.

REFERENCES

- Karumanchi SA, Maynard SE, Stillman IE, Epstein FH, Sukhatme VP. Preeclampsia: a renal perspective. *Kidney Int* 2005;67:2101-13.
- Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *Lancet* 2005;365:785-99.
- Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol* 2009;33:130-7.
- Fisher SJ. The placental problem: linking abnormal cytotrophoblast differentiation to the maternal symptoms of preeclampsia. *Reprod Biol Endocrinol* 2004;2:53.
- Kopcow HD, Karumanchi SA. Angiogenic factors and natural killer (NK) cells in the pathogenesis of preeclampsia. *J Reprod Immunol* 2007;76:23-9.
- Eastbrook G. The role of decidual natural killer cells in normal placentation and in the pathogenesis of preeclampsia. *Obstet Gynaecol Can* 2008;30:467-76.
- Kumar V, Medhi B. Emerging role of uterine natural killer cells in establishing pregnancy. *Iran J Immunol* 2008;5:71-81.
- Kopcow HD, Karumanchi SA. Angiogenic and natural killer (NK) cells in the pathogenesis of preeclampsia. *J Reprod Immunol* 2007;76:23-9.
- Milne F, Redman C, Walker J. The pre-eclampsia community guideline (PRECOG): how to screen for and detect onset of preeclampsia in the community. *Br Med J* 2005;330:576-80.
- Steegers EAP. Pre-eclampsia. *Lancet* 2010;376:631-44.
- Uzan Jennifer, Marie Carboneel, Olivier Piconne. Pre-eclampsia: pathophysiology, diagnosis, and management. *Vasc Health Risk Manage* 2011;7:467-74.
- Ng EH, Chan CC, Tang OS, Yeung WS, Ho PC. The role of endometrial and subendometrial vascular flow measured by three-dimensional power doppler ultrasound in the prediction of pregnancy during frozen-thawed embryo transfer cycles. *Hum Reprod* 2006;21:1612-7.
- Burton GJ, Jauniaux E. Placental oxidative stress: from miscarriage to preeclampsia. *J Soc Gynecol Investig* 2004;11:342-52.
- Jauniaux E, Watson AL, Hempstock J, Bao YP, Skepper JN, Burton GJ. Onset of maternal arterial blood flow and placental oxidative stress. A possible factor in human early pregnancy failure. *Am J Pathol* 2000;157:2111-22.
- Plasencia W, Maiz N, Bonino S, Kaihura C, Nicolaidis KH. Uterine artery Doppler at 11+0 to 13+6 w in the prediction of pre-eclampsia. *Ultrasound Obstet Gynecol* 2007;30:742-9.
- Burton GJ. Oxygen, the janus gas; its effects on human placental development and function. *J Anat* 2009;215:27-35.
- Burton GJ, Yung HW, Cindrova Davies T, Charnock Jones DS. Placental endoplasmic reticulum stress and oxidative stress in the pathophysiology of unexplained intrauterine growth restriction and early onset preeclampsia. *Placenta* 2009;30(Suppl A):43-8.
- Jauniaux E, Poston L, Burton GJ. Placental-related diseases of pregnancy: Involvement of oxidative stress and implications in human evolution. *Hum Reprod Update* 2006;12:747-55.
- Brosens JJ, Parker MG, McIndoe A, Pijnenborg R, Brosens IA. A role for menstruation in preconditioning the uterus for successful pregnancy. *Am J Obstet Gynecol* 2009;200:615.
- Huppertz B. Placental origins of preeclampsia: challenging the current hypothesis. *Hypertension* 2008;51:970-5.
- Myers J, Mires G, Macleod M, Baker P. In preeclampsia, the circulating factors capable of altering *in vitro* endothelial function precede clinical disease. *Hypertension* 2005;45:258-63.
- Irani RA, Xia Y. The functional role of the renin-angiotensin system in pregnancy and preeclampsia. *Placenta* 2008;29:763-71.

COMPARISON IMMUNOHISTOCHEMISTRY EXPRESSION OF DESIDUAL NATURAL KILLER (dNK) IN SEVERE PREECLAMPSIA AND NORMAL PREGNANCY

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

13%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	journals.lww.com Internet Source	5%
2	repository.unair.ac.id Internet Source	3%
3	S. Ananth Karumanchi, Phyllis August, Tiina Podymow. "Renal Complications in Normal Pregnancy", Elsevier BV, 2010 Publication	2%
4	Submitted to University of Sunderland Student Paper	1%
5	Tinnakorn Chaiworapongsa, Roberto Romero, Francesca Gotsch, Jimmy Espinoza et al. "Low maternal concentrations of soluble vascular endothelial growth factor receptor-2 in preeclampsia and small for gestational age", The Journal of Maternal-Fetal & Neonatal Medicine, 2009 Publication	1%

6	obgin.usu.ac.id Internet Source	1%
7	Charles J. Lockwood, S. Joseph Huang, Chie-Pein Chen, Yingqun Huang et al. "Decidual Cell Regulation of Natural Killer Cell–Recruiting Chemokines", <i>The American Journal of Pathology</i> , 2013 Publication	1%
8	"Proceedings of the Andalas International Public Health Conference 2017" , <i>BMC Public Health</i> , 2017 Publication	1%
9	www.pubmedcentral.nih.gov Internet Source	1%
10	talenta.usu.ac.id Internet Source	1%

Exclude quotes On
 Exclude bibliography On

Exclude matches < 1%